# **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 4:		(11) International Publicati n Number: WO 89/ 08403
A23L 1/30, 2/00	A1	43) International Publication Date: 21 September 1989 (21.09.89
(21) International Application Number: PCT/SE (22) International Filing Date: 9 March 1989 (31) Priority Application Number:	·	ropean patent), CH (European patent), DE (European patent), DK, FI, FR (European patent), GF (European patent), IT (European patent), IT (European patent), NO, SF
(32) Priority Date: 9 March 1988 (33) Priority Country:	•	Published  With international search report.  With amended claims.
(71)(72) Applicants and Inventors: MOLIN, Nils [SI Månsgatan 11 B, S-222 29 Lund (SE). Al SON, Carl-Erik [SE/SE]; Kristinavägen 40, Saltsjö-Duvnäs (SE). BENGMARK, Stig Box 5003, S-220 05 Lund (SE). LARSSON, I SE]; Norra Villavägen 7 B, S-237 00 Bjärred	LBERT S-131 [SE/SI Kåre [S	
(74) Agent: AWAPATENT AB; Box 5117, S-200 7 (SE).	1 Maln	

(54) Title: NUTRIENT COMPOSITION AND METHOD FOR THE PREPARATION THEREOF

### (57) Abstract

A nutrient composition for administration to patients in, for example, feeding by tube, or for use as a health drink is described. The nutrient composition comprises a fermented cereal-based product, enzyme and, optionally, further nutrient components, in combination with lactobacilli. Also described is a method of preparing a nutrient composition for administration to patients, inter alia in feeding by tube, or for use as a health drink, in which method a cereal flour is mixed with water,  $\alpha$ -amylase and, optionally, a protease, the mixture is brought to the boil under agitation, allowed to cool and mixed with  $\beta$ -glucanase which is allowed to act until the viscosity of the mixture has decreased to below 0.020 Pas, whereupon the mixture is fermented.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

	٠.				•
AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	П	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland .	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark	MC	Мопасо	ÜS	United States of America
स	Finland	MG	Madagagaga		

10

15

20

25

30

1

# NUTRIENT COMPOSITION AND METHOD FOR THE PREPARATION THEREOF

The present invention relates to a nutrient composition for administration to patients, inter alia for feeding by tube, and for use as a health drink, or as a nutrient for horses, as well as to a method for the preparation of such a nutrient composition.

One medical problem in major operations of the abdomen is the general collapse of the organs due to infections from the large intestine. One way of reducing the risks of infections is to feed the patient by tube, thereby to keep the intestine working. This method imposes special demands on the tube-administered nutrient. Not only must the nutrient be nutrimentally complete; it must also be so thin-bodied that it can be administered via a thin plastics catheter, without necessitating a daily supply of more than 2-3 liters of liquid.

Furthermore, it is a great advantage if the nutrient can affect the composition of the patient's intestinal flora which frequently is disturbed in newly operated patients, and this may sometimes result in symptoms of poisoning. This applies especially to patients who have been or are being treated with antibiotics.

A further field of application where there is a need for specific nutrients is horse-racing. Horses, under stress, for example in connection with a race, frequently contract diarrhoea, which may affect their performance.

Present-day nutrient compositions used for feeding by tube substantially consist of fractionated food components recombined according to the recommendations of dieticians. The known nutrient compositions suffer from a number of shortcomings. If they have a suitable low viscosity for administration through narrow tubes, their concentration of the requisite nutrients will

10

15

instead be too low, and therefore very large amounts must be given. Furthermore, the known nutrient compositions have a disagreeable taste and do not affect the patient's intestinal flora.

Indicia showing that certain intestine-specific lactobacilli have a special ability to counteract the establishment of undesired microorganisms in the intestine, have been reported by, for example, Wallström et al, 1987, J. Appl. Bacteriol. 62:513-520; Sherwood et al, 1987, Lancet, Dec. 26, 1519.

It is an object of the present invention to provide a nutrient composition for administration to patients, inter alia in feeding by tube, or for use as a health drink, said composition having a viscosity which is suitable for administration by tube, which is tasty, and which, furthermore, contains lactobacilli suitable for the intestine.

The nutrient composition according to the invention, especially when it contains fermented oats, is also highly suitable for administration to horses before a race for the prevention of diarrhoea.

A further object of the invention is to provide a method of preparing a nutrient composition for administration to patients, inter alia for feeding by tube, or for use as a health drink.

The nutrient composition according to the invention comprises a fermented cereal-based product, enzymes and, optionally, further nutrient components, in combination with lactobacilli.

The method according to the invention is characterised in that cereal flour is mixed with water and  $\alpha$ -amylase, and that the mixture is brought to the boil under agitation, allowed to cool and mixed with  $\beta$ -glucanase which is allowed to act until the viscosity of the mixture has decreased to below 0.020 Pas, whereupon the mixture is fermented.

A patient who is fed by tube can be supplied with the suitable lactobacilli in basically three

20

25

30

35

15

20

25

30

different ways, viz. (i) the desired lactobacillus can be utilised for fermenting the product; (ii) a lactobacillus culture can be added separately to the fermented product; and (iii) the lactobacillus culture and the rest of the nutrient composition can be administered separately.

The lactobacillus desired in the final product is preferably used already for fermenting the product, whereby there is conveniently obtained a fermented product containing the desired culture, without necessitating the addition of further bacteria cultures.

The lactobacilli supplied not only have a favourable effect on the patient's intestinal flora; they also contribute to reducing the viscocity of the nutrient composition during the fermentation process. Proteins are degraded to short dextrins and amino acids.  $\beta\text{-Glucanes}$  are degraded by the enzyme  $\beta\text{-glucanase}$ . The result is a readily digested product which is rich in flavour and has adequate shelf life.

In the preferred embodiment, the fermenting ability of the lactobacilli is utilised, and at the same time the patient's need of viable lactobacilli is satisfied.

The lactobacillus cultures selected are special cultures which in vitro have certain characteristics significant to the contemplated use. Suitable cultures are obtained by extracting and isolating lactobacilli that have established themselves in the intestine of healthy persons. The following characteristics are taken into account in selecting a suitable production strain:

- the ability to ferment oat-flour
- the specific growth rate
- the ability to rapidly reduce pH during the fermentation process
- 35 suitable final pH upon fermentation
  - the acid fermentation pattern from glucose

- survival upon freeze-drying
- resistance to bile
- resistance to antibiotics
- plasmide contents
- 5 the ability to spontaneously adhere to the intestine
  - reuterine production in the presence of glycerol
  - cholesterol interaction
  - the ability to produce desirable flavouring agents
  - the ability to degrade β-glucanes
- All of these characteristics need not occur in one and the same culture, but it is recommended to choose a culture that has as many as possible of these characteristics.
- A preferred culture is an isolate of intestine-specific Lactobacillus reuteri and a preparation of lipase
  enzymes stimulating the production of antimicrobial
  substances (reuterine) by said lactobacillus. Other
  useful organisms are, for example, Lactobacillus fermentum,
  L. acidofilus, L. alimentarius, L. brevis sp. lindneri,
- 20 L. plantarum, L. leuconostoc and L. dextranicum.

The basis of the nutrient composition is cereal flour. It has surprisingly been found that especially a suspension of oat-flour in water is an ideal basis for lactic acid fermentation. Of all cereals, oat

- has the best amino acid balance, and the lipid fraction contains the maximum amount of polar lipids. It has been shown, by works in progress, that this lipid type has a favourable effect on the gastric mucosa.
- In countries in which there is a shortage of oats, also other cereals, such as corn, may however be used.

In order to obtain a nutrient composition which is as complete as possible, the composition can be mixed with soya flour before fermentation to supplement the protein and fat contents.

Furthermore, minerals and vitamin contents may be supplemented before or after fermentation.

In the method, the cereal flour is suspended in water, and  $\alpha$ -amylase is added, either in the form of malt flour, pure  $\alpha$ -amylase, or in the form of  $\alpha$ -amylase-containing microorganisms. The suspension is brought to the boil under agitation to the suitable temperature which, if it is an oat suspension, is maximally 95°C. Then, the suspension is allowed to cool about 50°C, whereupon  $\beta$ -glucanase is added which is allowed to act for 1-2 hours at about 50°C, where-

upon the viscosity has decreased to about 0.020 Pas.

The cereal flour can also be treated with various proteases in order to degrade the proteins therein.

It is recommended that the product be then enriched with soya flour.

Finally, the mixture is mixed with a suitable bacteria culture, preferably the lactobacillus culture that is to be present in the final composition, and fermented at a temperature that should lie between 30 and 40°C.

The composition according to the invention may be used either as it is, in the form of an aqueous solution, or the aqueous solution can be concentrated and diluted immediately before use. Such a solution has an estimated shelf life of about 8 days at cooling temperature.

Alternatively, the nutrient solution can be freezedried and then dissolved in water before use. An additionally improved shelf life is obtained if the cereal flour, preferably the oat-flour is defatted before use by extraction in a solvent, such as ethanol or supercritical carbon dioxide, whereupon the resulting oil, after emulsifying and spray-drying is recycled to the product after freeze-drying.

The method described above thus gives a fermented

nutrient composition which is optimal for enteral

nutrition and which contains physiologically active

cells of a specially selected lactobacillus culture

10

in high concentration. Furthermore, the product gives a rapid colonisation in the small intestine.

The preferred freeze-drying imparts to the nutrient composition a much increased resistance to fat oxidation, whereby a satisfactory shelf life is obtained.

In two instances, the solution was administered to patients under intensive care for nutritional discorders and sepsis due to colonisation of aerobic intestinal bacteria. When the treatment began, both patients were apathetic because no previous treatment had proved successful. After administration of the nutrient solution according to the invention, containing lactobacilli, by tube to the intestine, both patients made a dramatic recovery.

The nutrient preparation according to the invention can be used also as a supplementary nourishment in general, for example in the form of a health drink.

The invention will now be described in more detail with reference to the following nonrestrictive Example.

## 20 EXAMPLE

35

34 g oat-flour (Mp oat-flour, product name) are suspended in 100 g water together with 3.4 g malt flour alternatively 1 g pure  $\alpha$ -amylase. The oat-flour suspension is brought to the boil under agitation to 95°C, whereupon it is allowed to cool to about 50°C. Then, 1%  $\beta$ -glucanase (calculated on the amount of oat-flour) is added and allowed to act for 1-2 hours at 50°C, whereby the viscosity is decreased to about 0.020 Pas.

After that, the suspension is enriched with soya flour.

Finally, a lactobacillus culture is added, selected with due regard to the characteristics mentioned in the general part of the specification, whereupon the suspension is fermented at, for example, 37°C for 20 hours.

-

To further enrich the product, further fat, sodium, vitamin A, vitamin D, riboflavin, vitamin B<sub>6</sub>, folic acid and ascorbic acid and, optionally, also potassium and calcium are added.

The product prepared in this manner has the energy and nutrient contents stated in Table 1.

TABLE 1

		No.	ENRICHMENT	
			Alt. 1	Alt. 2
Recommended		Oat sus-	Skim milk	Soya flour
contents/100 m	il	pension/100 ml	powder 3.7 g	2.8 g
Energy KJ	424	424 *	475 *	513 *
Protein g	4.7	3.4	4.7 *	4.7 *
Fat g	3.5	1.8	1.8	1.82
Carbohydr. g	11.8	16.6 *	18.5 *	17.6 *
Ca.mg	>51	13.5	58.9 *	20.8
Fe mg	>0.94	1.32 *	2.46 *	1.62 *
P mg	70	98 *	133 *	116 *
Mg mg	>28 g	35 *	40 *	44 *
Na mg	80	1.25	21	1.28
K mg	148	84	150 *	134
ret. eqv.mg >	0.05	*	0.011	0.0001
vit. D μg >	0.28	4 - 4	traces	-
Thiamine mg >	0.070	0.15 *	0.19 *	0.18 *
Ribofl. mg >			0.11 *	0.04
vit. B <sub>6</sub> mg >	0.11	0.04	0.053	0.059
Folic acid µg	40	0.006	0.006	0.06
Ascorb. mg >	3.3	·"·	0.3	-

<sup>\*</sup> Satisfies demands for recommended amount of nutrition

10

30

#### CLAIMS

- 1. Nutrient composition, c h a r a c t e r i s e d in that it comprises a fermented cereal-based product, enzymes and, optionally, further nutrient components, in combination with lactobacilli.
- 2. Nutrient composition as claimed in claim 1, c h a r a c t e r i s e d in that the cereal-based product is a product based on oat-flour.
- 3. Nutrient composition as claimed in claim l or 2, characterised in that the cereal-based product is fermented with said lactobacilli.
- 4. Nutrient composition as claimed in claim 1 or 2, character is ed in that the lactobacilli have been added to the composition after fermentation with another culture.
- 5. Nutrient composition as claimed in claim 1 or 2, character is ed in that the lactobacilli are present in a form separate from the remainder of the composition.
- 6. A nutrient composition as claimed in one or 20 more of the preceding claims, c h a r a c t e r i s e d in that the enzymes are  $\alpha$ -amylase,  $\beta$ -glucanase and, optionally, a protease.
- 7. Nutrient composition as claimed in one or more of the preceding claims, characterised 25 in that it comprises, as further nutrient components, soya flour and/or supplementary mineral substances and vitamins.
  - 8. Nutrient composition as claimed in one or more of the preceding claims, character is ed in that it is in the form of an aqueous solution or a freeze-dried powder.
    - 9. A method of preparing a nutrient composition, characterised in that
    - a cereal flour is mixed with water, α-amylase

and, optionally, a protease;

- the mixture is brought to the boil under agitation;
- is allowed to cool;
- is mixed with β-glucanase which is allowed to
   act until the viscosity of the mixture has decreased to below 0.020 Pas; and
  - is fermented.
- 10. A method as claimed in claim 9, c h a r a c t e r i s e d in that the cereal flour utilised is
  10 oat-flour.
  - 11. A method as claimed in claim 9 or 10, c h a r a c t e r i s e d in that the mixture is fermented
    with lactobacilli.
- 12. A method as claimed in claim 9 or 10, c h a 15 r a c t e r i s e d in that lactobacilli are admixed to the mixture after fermentation.
  - 13. A method as claimed in one or more of claims 9-12, characterised in that soya flour is admixed to the mixture before fermentation.
- 20 14. A method as claimed in one or more of claims 9-13, character is ed in that the mixture is enriched with supplementary mineral substances and vitamins before or after fermentation.
- 15. A method as claimed in one or more of claims 25 9-14, c h a r a c t e r i s e d in that the mixture is freeze-dried after fermentation.
  - 16. A method as claimed in claim 15, c h a r a c t e r i s e d in that the cereal flour is defatted before it is mixed with water and  $\alpha$ -amylase, and in that the removed fat after emulsifying and spray-drying is recycled to the freeze-dried composition.

### AMENDED CLAIMS

[received by the International Bureau on 20 July 1989 (20.07.89)

original claims 1-16 replaced by amended claims 1-11 (3 pages)]

- l. Nutrient composition which is optimal for enteral nutrition and which has a low viscosity, c h a r a c t e r i s e d in that it comprises an aqueous solution of a fermented product based on oat-flour,  $\alpha$ -amylase,  $\beta$ -glucanase and, optionally, a protease and, optionally, further nutrient components, in combination with viable lactobacilli which have the ability to spontaneously adhere to the human intestine and which are isolated from the human intestine.
- 2. Nutrient composition as claimed in claim 1, c h a r a c t e r i s e d in that the lactobacilli have been selected to have as many of the following characteristics as possible:
- the ability to ferment oat-flour
- the specific growth rate
- the ability to rapidly reduce pH during the fermentation process
- suitable final pH upon fermentation
- the acid fermentation pattern from glucose
- survival upon freeze-drying
- resistance to bile
- resistance to antibiotics
- plasmide contents
- reuterine production in the presence of glycerol
- cholesterol interaction
- the ability to produce desirable flavouring agents
- the ability to degrade β-glucanes.
- 3. Nutrient composition as claimed in claim 1 or 2, c h a r a c t e r i s e d in that the oat-flour based product is fermented with said lactobacilli.
- 4. Nutrient composition as claimed in claim 1 or 2, c h a r a c t e r i s e d in that the lacto-bacilli have been added to the composition after fermentation with another culture.

- 5. Nutrient composition as claimed in one or more of the preceding claims, c h a r a c t e r i s e d in that it comprises, as further components, soya flour and/or supplementary mineral substances and vitamins.
- 6. Nutrient composition as claimed in one or more of the preceding claims, characterised in that it is in the form of a freeze-dried powder dissolved in water.
- 7. A method of preparing a nutrient composition which is optimal for enteral nutrition and which has a viscosity of 0.020 Pas at most, character rised in that
- an oat-flour is mixed with water,  $\alpha$ -amylase and, optionally, a protease;
- the mixture is brought to the boil under agitation;
- is allowed to cool;
- is mixed with  $\beta$ -glucanase which is allowed to act until the viscosity of the mixture has decreased to below 0.020 Pas; and
- is fermented, preferably by using viable lactobacilli; which have the ability to spontaneous adhere to the human intestine and which have been isolated from the human intestine,
- or, alternatively, if not used for the fermentation, said viable lactobacilli are added after fermentation.
- 8. A method as claimed in claim 7, character is ed in that soya flour is admixed to the mixture before fermentation.
- 9. A method as claimed in claim 7 or 8, c h a r a c t e r i s e d in that the mixture is enriched with supplementary mineral substances and vitamins before or after fermentation.
- 10. A method as claimed in one or more of claims 7-9, characterised in that the mixture is freeze-dried after fermentation and then again dissolved in water before use.

11. A method as claimed in claim 10, c h a - r a c t e r i s e d in that the oat-flour is defatted before it is mixed with water and  $\alpha$ -amylase, and in that the removed fat after emulsifying and spray-drying is recycled to the freeze-dried composition.

### INTERNATIONAL SEARCH REPORT

International Application No PCT/SE89/00114

		F SUBJECT MATTER (if several class Patent Classification (IPC) or to both No		
		•	4	
	L 1/30	2/00		<del></del>
	9221131129	Minimum Docum	entation Searched 7	
Classification	on System		Classification Symbols	
IPC	i	•	305, 2/00, /38; A 2	3 C 9/133
US	C1	<u>126</u> :18,20,48,49,52,	330.3,590	
		Documentation Searched other to the Extent that such Document	than Minimum Documentation ts are included in the Fields Searched <sup>8</sup>	
SE,	NO, DK,	FI classes as abo	ve	
III. DOCU		IIDERED TO SE RELEVANT		
alegory •	Citation o	Document, 11 with Indication, where ap	propriate, of the relevant passages 12	Relevant to Claim No. 13
x	US, A,	4 056 637 (HAGIWA 1 November 1977 See claims 1-4	RA et al)	1,3,4,6,8,
x	US, A,	2 452 534 (G.A. J 2 November 1948 See claim 1	EFFREYS)	1,3
X	US, A,	2 194 672 (C.M. PO 26 March 1940 See claims 1,2	ORTER)	1,3,7
x	C-342,	Abstracts of Japan abstract of JP 60- 985-11-30	n, Vol 10, No 112, -241848,	1,3,9
x	C-343,	Abstracts of Japa abstracts of JP 60 985-12-07	an, Vol 10, No 120, D-248131,	1,3,9
х	C-51,	Abstracts of Japar abstract of JP 56-1 981-02-05		1,4,9
"A" docu cons "E" earlie filing "L" docu which citati "O" docu other	iment defining the idered to be of a comment but a date iment which many is cited to estion or other speciment referring to means.	ted documents: 16 is general state of the art which is not particular relevance published on or after the international or throw doubts on priority claim(s) or ablish the publication date of another tial reason (as specified) or noral disclosure, use, exhibition or prior to the international filing date but date claimed	"T" later document published after the or priority date and not in conflicted to understand the principle invention.  "X" document of particular relevant cannot be considered novel or involve an inventive step document of particular relevant cannot be considered to involve document is combined with one ments, such combination being on the art.	ct with the application but or theory underlying the ce; the claimed invention cannot be considered to ce; the claimed invention an inventive step when the or more other such docu- physious to a person skilled
	FICATION		Date of Mailine of this International Co.	arch Banert
	Actual Completi	on of the international Search	Date of Melling of this international Se	aren Report
	Searching Aut	nority		<u> </u>
	-	ent Office	Inga-Karin Peterss	les serve

ategory •	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
x	Patent Abstracts of Japan, Vol 8, No 212, C-244, abstracts of JP 59-98672 publ 1984-06-07	1,3,9
ŀ		
Ì		
	·	
		1
	•	
	•	

.